Nebraska High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN8: Percentage of students who rarely or never wore a bicycle helmet (among students who had ridden a bicycle during the 12 months before the survey)	84.8	1,141	1.24	(82.2 - 87.1)	1.4
QN9: Percentage of students who rarely or never wore a seat belt (when riding in a car driven by someone else)	11.3	1,670	1.20	(9.1 - 13.9)	2.4
QN10: Percentage of students who rode with a driver who had been drinking alcohol (in a car or other vehicle one or more times during the 30 days before the survey)	22.3	1,677	1.39	(19.7 - 25.2)	1.9
QN11: Percentage of students who drove when drinking alcohol (one or more times during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	10.1	1,156	1.40	(7.6 - 13.3)	2.5
QN12: Percentage of students who texted or e-mailed while driving a car or other vehicle (on at least 1 day during the 30 days before the survey, among students who had driven a car or other vehicle during the 30 days before the survey)	49.4	1,167	2.22	(45.0 - 53.8)	2.3
QN15: Percentage of students who carried a weapon on school property (such as a gun, knife, or club on at least 1 day during the 30 days before the survey)	8.1	1,685	0.95	(6.4 - 10.2)	2.0

*The average design effect across all items for this survey is 1.9.

Nebraska High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN16: Percentage of students who did not go to school because they felt unsafe at school or on their way to or from school (on at least 1 day during the 30 days before the survey)	6.2	1,683	0.93	(4.6 - 8.4)	2.5
QN17: Percentage of students who were threatened or injured with a weapon on school property (such as a gun, knife, or club one or more times during the 12 months before the survey)	7.1	1,659	0.83	(5.6 - 8.9)	1.7
QN18: Percentage of students who were in a physical fight (one or more times during the 12 months before the survey)	19.7	1,649	1.08	(17.7 - 22.0)	1.2
QN19: Percentage of students who were injured in a physical fight (one or more times during the 12 months before the survey; injuries had to be treated by a doctor or nurse)	2.8	1,663	0.57	(1.9 - 4.2)	2.0
QN20: Percentage of students who were in a physical fight on school property (one or more times during the 12 months before the survey)	5.5	1,651	0.62	(4.4 - 6.9)	1.2
QN21: Percentage of students who were ever physically forced to have sexual intercourse (when they did not want to)	8.3	1,538	0.80	(6.8 - 10.0)	1.3

*The average design effect across all items for this survey is 1.9.

Nebraska High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN22: Percentage of students who experienced physical dating violence (one or more times during the 12 months before the survey, including being hit, slammed into something, or injured with an object or weapon on purpose by someone they were dating or going out with among students who dated or went out with someone during the 12 months before the survey)	8.1	983	1.21	(6.0 - 10.8)	1.9
QN23: Percentage of students who experienced sexual dating violence (one or more times during the 12 months before the survey, including kissing, touching, or being physically forced to have sexual intercourse when they did not want to by someone they were dating or going out with among students who dated or went out with someone during the 12 months before the survey)	8.8	972	1.01	(7.0 - 11.1)	1.2
QN24: Percentage of students who were bullied on school property (during the 12 months before the survey)	26.3	1,675	1.28	(23.8 - 28.9)	1.4
QN25: Percentage of students who were electronically bullied (including being bullied through e-mail, chat rooms, instant messaging, websites, or texting during the 12 months before the survey)	18.9	1,671	1.27	(16.5 - 21.5)	1.7

*The average design effect across all items for this survey is 1.9.

Nebraska High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN26: Percentage of students who felt sad or hopeless (almost every day for 2 or more weeks in a row so that they stopped doing some usual activities during the 12 months before the survey)	24.1	1,515	1.52	(21.2 - 27.3)	1.9
QN27: Percentage of students who seriously considered attempting suicide (during the 12 months before the survey)	14.6	1,552	1.00	(12.8 - 16.8)	1.2
QN28: Percentage of students who made a plan about how they would attempt suicide (during the 12 months before the survey)	13.3	1,546	0.98	(11.5 - 15.4)	1.3
QN29: Percentage of students who attempted suicide (one or more times during the 12 months before the survey)	8.9	1,330	0.79	(7.4 - 10.5)	1.0
QN30: Percentage of students who attempted suicide that resulted in an injury, poisoning, or overdose that had to be treated by a doctor or nurse (during the 12 months before the survey)	3.3	1,334	0.49	(2.5 - 4.5)	1.0
QN31: Percentage of students who ever tried cigarette smoking (even one or two puffs)	31.4	1,626	2.01	(27.6 - 35.6)	3.1
QN32: Percentage of students who smoked a whole cigarette before age 13 years (for the first time)	8.8	1,612	1.02	(6.9 - 11.0)	2.1
QN33: Percentage of students who currently smoked cigarettes (on at least 1 day during the 30 days before the survey)	13.3	1,606	1.54	(10.5 - 16.7)	3.3

*The average design effect across all items for this survey is 1.9.

Nebraska High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN34: Percentage of students who smoked more than 10 cigarettes per day (during the 30 days before the survey among students who currently smoked cigarettes on the days they smoked)	8.0	197	2.15	(4.6 - 13.4)	1.2
QN35: Percentage of students who usually obtained their own cigarettes by buying them in a store or gas station (during the 30 days before the survey among students who currently smoked cigarettes and who were aged <18 years)	9.7	169	2.35	(6.0 - 15.5)	1.1
QN36: Percentage of students who tried to quit smoking cigarettes (among students who currently smoked cigarettes during the 12 months before the survey)	48.7	186	4.11	(40.6 - 56.9)	1.3
QN37: Percentage of students who currently used smokeless tobacco (chewing tobacco, snuff, or dip on at least 1 day during the 30 days before the survey)	9.3	1,671	1.06	(7.4 - 11.7)	2.2
QN38: Percentage of students who currently smoked cigars (cigars, cigarillos, or little cigars on at least 1 day during the 30 days before the survey)	8.1	1,667	0.99	(6.3 - 10.3)	2.2
QN39: Percentage of students who ever used electronic vapor products (e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens such as blu, NJOY, or Starbuzz)	38.2	1,624	1.85	(34.6 - 41.9)	2.4

*The average design effect across all items for this survey is 1.9.

Nebraska High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN40: Percentage of students who currently used electronic vapor products (e-cigarettes, e-cigars, e-pipes, vape pipes, vaping pens, e-hookahs, and hookah pens such as blu, NJOY, or Starbuzz on at least 1 day during the 30 days before the survey)	22.3	1,632	1.58	(19.3 - 25.6)	2.3
QN41: Percentage of students who ever drank alcohol (at least one drink of alcohol on at least 1 day during their life)	51.7	1,629	1.67	(48.4 - 55.0)	1.8
QN42: Percentage of students who drank alcohol before age 13 years (for the first time other than a few sips)	13.7	1,659	1.16	(11.5 - 16.1)	1.9
QN43: Percentage of students who currently drank alcohol (at least one drink of alcohol on at least 1 day during the 30 days before the survey)	22.7	1,572	1.65	(19.6 - 26.2)	2.5
QN44: Percentage of students who drank five or more drinks of alcohol in a row (within a couple of hours on at least 1 day during the 30 days before the survey)	14.3	1,627	1.60	(11.4 - 17.8)	3.4
QN45: Percentage of students who reported that the largest number of drinks they had in a row was 10 or more (within a couple of hours during the 30 days before the survey)	3.3	1,580	0.64	(2.3 - 4.9)	2.0

*The average design effect across all items for this survey is 1.9.

Nebraska High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN46: Percentage of students who usually obtained the alcohol they drank by someone giving it to them (among students who currently drank alcohol)	39.6	339	2.72	(34.3 - 45.1)	1.0
QN47: Percentage of students who ever used marijuana (one or more times during their life)	26.6	1,613	1.93	(22.9 - 30.6)	3.1
QN48: Percentage of students who tried marijuana before age 13 years (for the first time)	6.3	1,617	1.02	(4.5 - 8.6)	2.9
QN49: Percentage of students who currently used marijuana (one or more times during the 30 days before the survey)	13.7	1,638	1.60	(10.8 - 17.2)	3.5
QN50: Percentage of students who ever used cocaine (any form of cocaine, such as powder, crack, or freebase, one or more times during their life)	5.3	1,626	0.78	(4.0 - 7.1)	2.0
QN51: Percentage of students who ever used inhalants (sniffed glue, breathed the contents of aerosol spray cans, or inhaled any paints or sprays to get high, one or more times during their life)	8.1	1,645	0.86	(6.6 - 10.0)	1.6
QN52: Percentage of students who ever used heroin (also called "smack," "junk," or "China white," one or more times during their life)	2.5	1,641	0.70	(1.4 - 4.3)	3.3

*The average design effect across all items for this survey is 1.9.

Nebraska High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN53: Percentage of students who ever used methamphetamines (also called "speed," "crystal," "crank," or "ice," one or more times during their life)	4.2	1,612	0.67	(3.1 - 5.8)	1.8
QN54: Percentage of students who ever used ecstasy (also called "MDMA," one or more times during their life)	5.1	1,641	0.72	(3.9 - 6.8)	1.7
QN55: Percentage of students who ever used synthetic marijuana (also called "K2", "Spice", "fake weed", "King Kong", "Yucatan Fire", "Skunk", or "Moon Rocks", one or more times during their life)	7.5	1,635	1.08	(5.6 - 9.9)	2.8
QN56: Percentage of students who ever took steroids without a doctor's prescription (pills or shots, one or more times during their life)	3.5	1,644	0.65	(2.4 - 5.0)	2.0
QN57: Percentage of students who ever took prescription drugs without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, one or more times during their life)	13.5	1,643	1.34	(11.1 - 16.4)	2.5
QN58: Percentage of students who ever injected any illegal drug (used a needle to inject any illegal drug into their body one or more times during their life)	4.0	1,638	0.75	(2.7 - 5.8)	2.4

*The average design effect across all items for this survey is 1.9.

Nebraska High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN59: Percentage of students who were offered, sold, or given an illegal drug on school property (during the 12 months before the survey)	19.9	1,629	1.57	(16.9 - 23.2)	2.5
QN60: Percentage of students who ever had sexual intercourse	32.5	1,365	2.00	(28.6 - 36.6)	2.5
QN61: Percentage of students who had sexual intercourse before age 13 years (for the first time)	3.3	1,352	0.61	(2.3 - 4.8)	1.6
QN62: Percentage of students who had sexual intercourse with four or more persons (during their life)	8.0	1,360	0.94	(6.3 - 10.1)	1.6
QN63: Percentage of students who were currently sexually active (sexual intercourse with at least one person during the 3 months before the survey)	24.9	1,359	1.85	(21.4 - 28.7)	2.5
QN64: Percentage of students who drank alcohol or used drugs before last sexual intercourse (among students who were currently sexually active)	17.9	314	3.24	(12.4 - 25.3)	2.2
QN65: Percentage of students who used a condom (during last sexual intercourse among students who were currently sexually active)	57.0	314	3.49	(49.9 - 63.8)	1.6
QN66: Percentage of students who used birth control pills (before last sexual intercourse to prevent pregnancy among students who were currently sexually active)	20.7	308	2.46	(16.2 - 26.0)	1.1

*The average design effect across all items for this survey is 1.9.

Nebraska High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN69: Percentage of students who described themselves as slightly or very overweight	30.3	1,612	1.25	(27.9 - 32.9)	1.2
QN70: Percentage of students who were trying to lose weight	43.2	1,620	1.59	(40.1 - 46.4)	1.7
QN71: Percentage of students who did not drink fruit juice (100% fruit juices one or more times during the 7 days before the survey)	29.9	1,608	1.33	(27.3 - 32.6)	1.4
QN72: Percentage of students who did not eat fruit (one or more times during the 7 days before the survey)	10.3	1,596	1.01	(8.5 - 12.5)	1.8
QN73: Percentage of students who did not eat salad (during the 7 days before the survey)	39.4	1,613	1.29	(36.9 - 42.1)	1.1
QN74: Percentage of students who did not eat potatoes (during the 7 days before the survey)	30.6	1,607	1.33	(28.0 - 33.4)	1.3
QN75: Percentage of students who did not eat carrots (during the 7 days before the survey)	50.0	1,602	1.63	(46.8 - 53.3)	1.7
QN76: Percentage of students who did not eat other vegetables (during the 7 days before the survey)	19.1	1,606	1.17	(16.8 - 21.5)	1.4
QN77: Percentage of students who did not drink a can, bottle, or glass of soda or pop (not including diet soda or diet pop, during the 7 days before the survey)	24.3	1,600	1.23	(22.0 - 26.9)	1.3

*The average design effect across all items for this survey is 1.9.

Nebraska High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN78: Percentage of students who did not drink milk (during the 7 days before the survey)	17.3	1,588	1.25	(14.9 - 19.9)	1.7
QN79: Percentage of students who did not eat breakfast (during the 7 days before the survey)	13.3	1,587	1.02	(11.4 - 15.4)	1.4
QN80: Percentage of students who were physically active at least 60 minutes per day on 5 or more days (doing any kind of physical activity that increased their heart rate and made them breathe hard some of the time during the 7 days before the survey)	52.8	1,584	1.98	(48.8 - 56.7)	2.5
QN81: Percentage of students who watched television 3 or more hours per day (on an average school day)	20.1	1,588	1.11	(18.0 - 22.4)	1.2
QN82: Percentage of students who played video or computer games or used a computer 3 or more hours per day (for something that was not school work on an average school day)	31.5	1,580	1.63	(28.4 - 34.9)	1.9
QN83: Percentage of students who attended physical education classes on 1 or more days (in an average week when they were in school)	49.0	1,565	1.95	(45.1 - 52.9)	2.4
QN84: Percentage of students who played on at least one sports team (run by their school or community groups during the 12 months before the survey)	64.3	1,566	1.65	(60.9 - 67.5)	1.9

*The average design effect across all items for this survey is 1.9.

Nebraska High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN85: Percentage of students who were ever tested for HIV (not including tests done when donating blood)	9.3	1,438	0.78	(7.8 - 11.0)	1.1
QN86: Percentage of students who saw a dentist (for a check-up, exam, teeth cleaning, or other dental work during the 12 months before the survey)	75.2	1,457	1.38	(72.4 - 77.8)	1.5
QN87: Percentage of students who had ever been told by a doctor or nurse that they had asthma	20.0	1,445	1.31	(17.5 - 22.8)	1.5
QN88: Percentage of students who had 8 or more hours of sleep (on an average school night)	31.9	1,452	1.74	(28.6 - 35.5)	2.0
QN89: Percentage of students who made mostly A's or B's in school (during the 12 months before the survey)	79.4	1,447	1.67	(75.9 - 82.5)	2.5
QN90: Percentage of students who talked on a cell phone while driving (on at least 1 day during the 30 days before the survey, among students who drove a car or other vehicle)	53.8	1,235	2.17	(49.5 - 58.1)	2.3
QN91: Percentage of students who reported liquor as the type of alcohol they usually drank (such as vodka, rum, scotch, bourbon, or whiskey, during the 30 days before the survey, among students who drank alcohol and who have a usual type of alcohol that they drink)	39.8	319	3.14	(33.7 - 46.2)	1.3

*The average design effect across all items for this survey is 1.9.

Nebraska High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN92: Percentage of students who reported their home or another person's home as the place they usually drank alcohol (during the 30 days before the survey, among students who drank alcohol)	84.5	375	3.14	(77.1 - 89.7)	2.8
QN93: Percentage of students who usually used marijuana by smoking it in a joint, bong, pipe, or blunt (during the 30 days before the survey, among students who used marijuana)	87.7	231	2.66	(81.3 - 92.1)	1.5
QN94: Percentage of students who currently took a prescription drug without a doctor's prescription (such as OxyContin, Percocet, Vicodin, codeine, Adderall, Ritalin, or Xanax, one or more times during the 30 days before the survey)	8.1	1,644	0.91	(6.5 - 10.1)	1.8
QN95: Percentage of students who went without eating for 24 hours or more to lose weight or to keep from gaining weight (also called fasting, during the 30 days before the survey)	12.6	1,603	1.14	(10.4 - 15.0)	1.9
QN96: Percentage of students who took some diet pills, powders, or liquids without a doctor's advice to lose weight or to keep from gaining weight (during the 30 days before the survey)	6.7	1,592	0.76	(5.3 - 8.4)	1.5
QN97: Percentage of students who vomited or took laxatives to lose weight or to keep from gaining weight (during the 30 days before the survey)	6.3	1,592	0.71	(5.0 - 7.8)	1.4

*The average design effect across all items for this survey is 1.9.

Nebraska High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN98: Percentage of students who did not drink a can, bottle, or glass of a sports drink (such as Gatorade or PowerAde, not including low-calorie sports drinks such as Propel or G2, during the 7 days before the survey)	34.2	1,599	1.50	(31.2 - 37.2)	1.6
QN99: Percentage of students who drank a can, bottle, or glass of a sugar-sweetened beverage (such as lemonade, sweetened tea or coffee drinks, flavored milk, Snapple, Sunny Delight, or energy drinks such as Red Bull, not including soda or pop, sports drinks, energy drinks, or 100% fruit juice, one or more times per day during the 7 days before the survey)	13.0	1,604	0.81	(11.5 - 14.7)	0.9
QN100: Percentage of students who have to avoid some foods because eating the food could cause an allergic reaction (such as skin rashes, swelling, itching, vomiting, coughing, or trouble breathing)	12.8	1,559	1.12	(10.7 - 15.2)	1.7
QN101: Percentage of students who did exercises to strengthen or tone their muscles (such as push-ups, sit-ups, or weight lifting, on three or more days during the 7 days before the survey)	54.3	1,586	1.77	(50.8 - 57.8)	2.0
QN102: Percentage of students who spend more than 20 minutes actually exercising or playing sports (during an average physical education (PE) class)	46.5	1,584	1.91	(42.8 - 50.4)	2.3

*The average design effect across all items for this survey is 1.9.

Nebraska High School Survey

Sample Statistics Report

Health Risk Behavior	Percentage	Number of Students	Standard Error	95% Confidence Interval	Design Effect*
QN103: Percentage of students who used an indoor tanning device (such as a sunlamp, sunbed, or tanning booth [not including getting a spray-on tan] one or more times during the 12 months before the survey)	14.2	1,459	1.41	(11.6 - 17.2)	2.4

*The average design effect across all items for this survey is 1.9.